

Parallel processing of visual space by neighboring neurons in mouse visual cortex

Spencer L Smith & Michael Häusser

Visual cortex shows smooth retinotopic organization on the macroscopic scale, but it is unknown how receptive fields are organized at the level of neighboring neurons. This information is crucial for discriminating among models of visual cortex. We used *in vivo* two-photon calcium imaging to independently map ON and OFF receptive field subregions of local populations of layer 2/3 neurons in mouse visual cortex. Receptive field subregions were often precisely shared among neighboring neurons. Furthermore, large subregions seem to be assembled from multiple smaller, non-overlapping subregions of other neurons in the same local population. These experiments provide, to our knowledge, the first characterization of the diversity of receptive fields in a dense local network of visual cortex and reveal elementary units of receptive field organization. Our results suggest that a limited pool of afferent receptive fields is available to a local population of neurons and reveal new organizational principles for the neural circuitry of the mouse visual cortex.

The organization of receptive fields (RFs) in the visual cortex¹ has been the subject of renewed and intense interest with the advent of new imaging and recording methods. A smooth retinotopic organization on the macroscopic scale has been demonstrated in mice using intrinsic signal optical imaging^{2,3}, corroborating earlier extracellular recordings^{4,5}. However, it remains unknown how RFs are organized on the microscopic scale, at the level of local networks of neighboring neurons. In monkeys, which also show smooth retinotopic organization, the scatter of RFs for units recorded simultaneously on the same electrode or during the same perpendicular penetration is quite large, approximately equal to the RF size⁶. More recent studies have confirmed that when multiple neurons are recorded simultaneously on the same electrode, they can have very different RFs^{7,8} and responses to natural stimuli⁹. However, these studies could only sample a small number of neurons that could fortuitously be recorded on a single electrode and discriminated. To determine the diversity of RFs and identify any underlying structure, it is necessary to systematically map larger samples in a defined local population.

Exploring the diversity of RFs within a local population can yield important insights into the function of visual cortex. The geometry of retinal ganglion cell mosaics¹⁰ can provide a constraint for the RFs of neurons in visual cortex^{10–12}. If neurons sample from a small pool of unique afferent RFs, then local populations of cortical neurons should show substantial redundancy in representing a few RFs. Alternatively, an ample pool of afferent RFs and a large amount of convergence could permit neighboring cortical neurons to integrate many different unique sets of afferents, thus yielding highly variable RFs and a low level of redundancy.

To address these issues, we used two-photon calcium imaging *in vivo* to obtain, to our knowledge, the first high-resolution RF maps for large numbers of neurons in a local population. By using

sparse-noise visual stimulation, we independently mapped the ON and OFF subregions of multiple neighboring neurons in mouse visual cortex. We discovered new features of RF micro-organization in visual cortex: shared and spanned receptive fields. These results can be used to constrain models of visual cortex, thus deepening our understanding of neocortical circuitry.

RESULTS

Mapping RFs for populations of neurons in visual cortex

We used bolus loading of calcium indicator dyes and two-photon microscopy^{13,14} to monitor spiking activity in a local population of layer 2/3 neurons spanning a square area (230 μm to a side) in mouse primary visual cortex during sparse-noise visual stimulation (Fig. 1a,b). A deconvolution-based algorithm was used to convert calcium signals into estimated spike rates. To calibrate the algorithm, we made loose cell-attached patch-clamp recordings from identified neurons in the population during two-photon calcium imaging and stimulus presentation (Fig. 1c). After calibrating using the electrophysiological data, the algorithm yielded spike-rate estimates that were highly correlated to the recorded spike trains ($r = 0.81 \pm 0.02$, $n = 19$; Fig. 1d), and the false positive rate was low ($4.4 \pm 1.7\%$).

The estimated spike rates were used to construct a spike-triggered average (Fig. 1e) for each neuron in the population. Visual stimuli consisted of random black and white dots on a gray background¹⁵. Using the estimated spike-rate time courses, we generated two spike-triggered averages, one for the white dots (the ON map) and one for the black dots (the OFF map). Maps were then smoothed using a Gaussian kernel equal in size to the smallest dot used in the stimulus set (1.3°), z-scored (normalized to units of standard deviations from the mean; Fig. 1f) and thresholded (Fig. 1g). With this method, we obtained maps for $44 \pm 4\%$ of imaged neurons (189/434 neurons in

Wolfson Institute for Biomedical Research and Department of Neuroscience, Physiology and Pharmacology, University College London, London, UK. Correspondence should be addressed to S.L.S. (spencer.smith@wibr.ucl.ac.uk).

Received 14 April; accepted 21 July; published online 15 August 2010; doi:10.1038/nn.2620




Figure 1 Mapping receptive fields with population calcium imaging and sparse-noise visual stimuli. **(a,b)** Presentation of sparse-noise visual stimuli **(a)** to a mouse during simultaneous population calcium imaging in visual cortex **(b)**. **(c)** Deconvolution of the calcium signals using parameters obtained from electrophysiology to obtain estimated spike rates. Micrograph scale bars in **b,c**, 20 μm ; vertical scale bars in **c**, 20% change in fluorescence, normalized to baseline ($\Delta F/F$) (top), 1 mV (middle), 2 spikes per frame (bottom). **(d)** Left, the mean correlation coefficient between deconvolved calcium signals and spike rates obtained from simultaneous on-cell recordings was 0.81 ± 0.02 . Right, detection reliability as a function of the number of spikes within one frame. Error bars, s.e.m. **(e)** Computation of a triggered average of stimulus frames, using the deconvolved calcium signals as an estimate of spike rate. Separate ON and OFF maps were generated using the white dots (top) and black dots (bottom). **(f)** Maps were filtered using a Gaussian kernel and z-scored using an area of the triggered average away from the RF. **(g)** These z-scored maps were thresholded to obtain RFs.

six mice). RFs for at least 20 neurons (range, 21–68) were obtained in each mouse. The relative ratio of ON and OFF subregions within a local population of neurons varied noticeably among individuals, but in contrast to that in macaque visual cortex, which is dominated by OFF responses¹⁶, overall did not show a consistent bias (mean, 1.08 ± 0.31 ; range, 0.39 to 2.17).

RF subregions varied in size (**Fig. 2a**) and were often elongated. To quantify RF subregion shape, we fit the subregions with ellipses and computed the half-short axis, which was $6.9 \pm 1.5^\circ$ (mean $\pm$ s.d., $n = 189$ RFs from six mice; **Fig. 2b**). This is similar to that of RF subregions of individual neurons in previous studies of mouse visual cortex obtained using electrophysiological techniques, which generally reported a 5–7° range for RF radius^{17–19}. The half-long axis of the RF subregions we measured was on average $11.1 \pm 4.3^\circ$ (mean $\pm$ s.d.; **Fig. 2c**); the difference between the two axis measurements reflects the often elongated structure of these RFs. The mean aspect ratio of the elliptical fits was 1.7 ± 0.6 (mean $\pm$ s.d.; **Fig. 2d**). The mean area of the RF subregions was $237 \pm 157^\circ$ (mean $\pm$ s.d.; **Fig. 2e**). ON and OFF subregions in individual RFs showed low overlap (**Supplementary Fig. 1**), indicating that most cells were simple cells, consistent with extracellular recordings from mouse V1 (refs. 18,20).

Overlap analysis of RF subregions

We next examined ensembles of subregions. Ensembles are defined here as all the subregions of a single sign (ON or OFF) from a

single population of neurons (20.7 ± 4.8 subregions per ensemble, $n = 6$ mice). Ensembles of subregions showed a scatter of $4.9 \pm 0.8^\circ$ ($n = 11$ ensembles from six mice; **Fig. 3a**). This scatter is comparable to the subregion size, consistent with findings in primates⁶. We then examined the degree of overlap in ensembles of subregions. To test whether the amount of overlap observed in the data could result from random positioning, we used a bootstrap method. For each ensemble of subregions, we constructed 1,000 new ensembles. These new ensembles used exactly the same population of subregions (no rotations or other distortions) but repositioned them randomly, preserving the subregion scatter observed in the original data (observed, $4.9 \pm 0.8^\circ$; shuffled, $4.9 \pm 0.4^\circ$; mean $\pm$ s.e.m.; $P = 0.99$, Wilcoxon signed-rank test; **Fig. 3a**). Thus, only the relative positions of subregions were randomized.

Next, we quantified the pairwise overlap of subregions using a simple metric: the proportion of the smaller of two subregions that overlapped the larger subregion. This metric ranges from 0 (indicating no overlap) to 1 (indicating that one subregion is completely contained within the other subregion). A high proportion of RF




Figure 2 Receptive field subregions obtained with population calcium imaging. **(a)** A representative set of examples, illustrating the diversity of subregion sizes and shapes, taken from eight different neurons across three different mice. For each example subregion, the thresholded, z-scored map (left; color scale as in **Fig. 1g**) and the subregion outline used in subsequent analysis (right) are displayed next to each other. The third and sixth subregions are OFF subregions; the rest, ON subregions. **(b–e)** Plots of the distributions of various geometric parameters for all subregions mapped ($n = 228$ subregions in six mice). **(b)** The half short axis length. **(c)** Half long axis length. **(d)** Aspect ratio of elliptical fits to observed subregions. **(e)** Distribution of areas of RF subregions. Arrows indicate the mean of each histogram.




Figure 3 Pairwise subregion overlaps are higher than expected for random positioning. **(a)** Left, multiple (12–15) overlaid subregions (top, ON; middle, OFF; bottom, OFF) in three different mice. For each point in space, the number of overlapping subregions is coded by color. Scatter is defined as the average deviation of each subregion center from the mean subregion center. Right, the subregions of each ensemble randomly repositioned three times. The random repositioning algorithm was designed to not change the scatter. Note that the random repositioning does not markedly alter the appearance of the ensembles. **(b)** Pairwise analysis, however, reveals that subregions overlapped more in their observed positions than when randomly repositioned (error bars, s.d.; $P < 10^{-5}$, Kolmogorov-Smirnov test). **(c)** Example of apparently shared subregions (left) and clustered subregion centers (right) in visual space. The subregion outlines and center markers have been color coded by hand for clarity. The two gray subregion center markers on the right represent subregions that do not seem to be shared with the green and blue groups.

subregion pairs showed a high degree of overlap (that is, an overlap index close to 1; **Fig. 3b**). To test whether this unexpectedly high degree of overlap was due to chance, we used the same measure for the randomly repositioned subregions (as shown in **Fig. 3a**). The randomly repositioned subregions showed substantially fewer incidents of high overlap ($P < 10^{-5}$, two-sample Kolmogorov-Smirnov test, **Fig. 3b**), indicating that the observed substantial overlap between RF subregions is a key, non-random feature of the organization of receptive fields in the mouse visual cortex. The distribution of full RFs (ON plus OFF subregions) was similar to that of repositioned subregions and was again significantly different from the subregions as observed ($P < 10^{-5}$, two-sample Kolmogorov-Smirnov test, **Supplementary Fig. 2**). Manual inspection of small ensembles of overlapping subregions revealed an apparent trend for subregions to be shared among multiple neurons (**Fig. 3c**). Therefore, we explicitly tested for the existence of shared subregions.


Shared RF subregions

We defined neurons with shared RF subregions as those whose subregions matched in shape, position and sign (ON or OFF). Such shared RF subregions (**Fig. 4a**) were observed frequently in all mice and were often shared across several (up to seven) neurons. When subregions were repositioned randomly in visual space while preserving scatter (see **Fig. 3a**), the incidence of shared subregions was significantly reduced ($P = 0.00021$, paired t -test, $n = 11$ subregion ensembles from six mice; **Fig. 4b**).

Figure 4 Features of local receptive field organization: shared subregions and spanned subregions. **(a)** Left, four examples showing subregions of individual (color-coded) neurons that are shared. Right, the relative position of the same neurons in visual cortex (same color code; the region of interest used for analysis is highlighted by a circle for ease of identification). **(b)** The number of shared subregions in the population decreases when the subregions are randomly repositioned. Each point represents one ensemble of subregions. **(c)** In the case of spanned subregions, the subregion of one neuron overlaps two or more non-overlapping subregions from other neurons. Four examples are shown. The upward (U) and temporal (T) directions in visual space, and the medial (M) and rostral (R) directions along the cortex, are indicated by arrows. **(d)** Again, the incidence of this spatial arrangement decreases when the subregions are randomly repositioned.

We then determined whether, in addition to sharing an area of visual space, neurons also shared the spatial structure of the spike-triggered average within the area. We computed the image correlation of the z -scored maps and the thresholded RF subregions. We found that the z -scored maps also showed high degrees of correlation, consistently higher than that of the thresholded RF subregions. This is further evidence of shared RF subregions (**Supplementary Fig. 3**).

What is the spatial relationship of neurons in visual cortex that share RF subregions? Two-photon calcium imaging of the neuronal population provides us with the exact location of all neurons recorded, allowing us to link their functional properties with their relative locations in the visual cortex. There was only a weak relationship between the distance separating individual neurons in the cortex and the overlap of their subregions in visual space (**Supplementary Fig. 4**). Therefore, although there is gross retinotopy on the macroscopic scale^{3,4}, visual cortex did not show smooth retinotopy on the scale of neighboring neurons.






Figure 5 ON and OFF subregions are offset with respect to each other in visual space. **(a)** The ON subregions in each mouse tend to cluster in an area of visual space spatially segregated from the OFF subregions, which also cluster together. Data from two mice are shown, with overlap density represented by amount of white (example 1 consists of 15 ON subregions and 18 OFF subregions; example 2 consists of 19 ON subregions and 15 OFF subregions). The upward (U) and temporal (T) directions in visual space are indicated by arrows. **(b)** When the subregions are randomly reassigned as ON or OFF subregions, the spatial segregation is lost. The observed clustering was decreased by randomly reassigning subregions as ON or OFF ($P < 0.02$, bootstrap method, $n = 4$ mice).

Correlations among neurons with shared RF subregions

The next question we addressed was whether neurons with shared RF subregions also show correlated activity. We computed activity correlations from the deconvolution-based spike-rate estimates rather than from the raw calcium imaging signals themselves to avoid overestimating the correlation coefficients (Supplementary Figs. 5 and 6). Although some neurons with shared RF subregions showed correlated activity, in general, neurons that shared RF subregions showed low correlations (Supplementary Fig. 5a–d), only slightly higher than that of the general population (correlation among neurons with shared RF subregions, 0.12 ± 0.12 , mean $\pm$ s.d., $n = 149$ pairs; correlation among neurons not sharing an RF subregion, 0.085 ± 0.088 , mean $\pm$ s.d., $n = 6,696$ pairs; $P = 0.0023$, Wilcoxon rank-sum test). Overall there was only a weak relationship between the correlation coefficient and the subregion overlap, as quantified by the overlap index (Pearson's $r = 0.13$; $P < 10^{-5}$; Supplementary Fig. 5f). There was a similarly weak relationship between the strength of the cross-correlation and the distance between cell bodies in visual cortex (Pearson's $r = -0.053$; $P = 0.0021$; Supplementary Fig. 5g). Therefore, even when the subregions of two neurons were identical (Supplementary Fig. 5b), individual neurons often responded to different visual stimuli (Supplementary Fig. 5e).

When we analyzed pairs of cells that both showed ON and OFF RF subregions and shared at least one RF subregion, we found that none (0/13) of these cell pairs shared both ON and OFF subregions. To further explore this difference, we examined the overlap and found that among these neurons, their subregions of the opposite sign overlapped about 30% less than did the shared subregions ($P = 0.0097$; Supplementary Fig. 6). This tendency for neurons to not share RF subregions of both signs simultaneously could contribute to the low correlations we observe. Further potential contributors to the low correlations include a low response reliability, differing selectivities to higher-order aspects of the stimulus not reflected in the spike-triggered average and active decorrelation due to intracortical circuitry²¹.

Spanned RF subregions

We also observed evidence for neurons integrating subregions into a single, larger subregion. We refer to this feature as 'spanned subregions': the subregion of one neuron overlaps two or more non-overlapping subregions of the same sign from other neurons (Fig. 4c). Spanned subregions were observed in all mice but not all ensembles. When subregions were repositioned randomly in visual space while preserving

scatter, the incidence of this feature was significantly reduced ($P = 0.0234$, paired Wilcoxon signed-rank test, $n = 8$ subregion ensembles from six mice; Fig. 4d). This indicates that subregions that are not overlapping may themselves be encompassed by a larger spanning subregion that integrates them both.

Spatial offset of ON and OFF RF subregion ensembles

Having determined that subregions of the same sign frequently overlapped and thus formed clusters in visual space, we next sought to determine the relative relationships between clusters of ON and OFF subregions. To address this issue, the ensemble of subregions for a single local population of neurons was plotted in visual space and color coded to discriminate ON and OFF subregions. This revealed a clear spatial segregation of ON and OFF subregion populations (Fig. 5a). When subregions were randomly reassigned as ON or OFF subregions (while still maintaining the same total number of ON and OFF subregions), this segregation was not observed ($P < 0.02$, bootstrap, $n = 4$ mice; Fig. 5b). Thus, local populations of neurons encode light and dark stimuli in parallel from separate regions of visual space.

DISCUSSION

We have mapped RFs in local populations of layer 2/3 neurons in mouse visual cortex using calcium imaging combined with sparse-noise visual stimulation. This yielded a cellular-level view of a cortical retinotopic map, revealing independently sampled ON and OFF subregions for individual neurons mapped at high resolution. Within these local populations we found that, although local retinotopy was not smooth, neurons often showed highly similar RF subregions, precisely overlapping in visual space and sharing similar spatial structure. Moreover, large subregions often seemed to be composed of two or more smaller, non-overlapping RF subregions shown by other neurons in the same local population. Together, these results suggest that small RF subregions represent elementary units that are shared and integrated by local populations of neurons. This finding indicates that a small pool of afferents strongly influences the organization of RFs in local populations and places constraints on the functional organization of cortical circuitry^{10–12,22}.

Population imaging for dense sampling of receptive fields

This study provides the first characterization, to our knowledge, of the diversity of spatial RFs in local networks of the mammalian visual cortex. Previous studies of visual cortical activity using population calcium imaging used comparatively small numbers of stimuli to reveal the cellular-level cortical organization of orientation selectivity^{14,23} and ocular dominance^{24,25}. In our experiments, by combining sparse-noise visual stimuli with high-speed imaging, we were able to present thousands of visual stimuli and obtain high-resolution RFs from a large fraction of neurons in a local population simultaneously.

Such dense sampling is important for constructing accurate models of cortical function^{15,26}. In particular, our results would have been difficult to obtain using smaller samples of each local population, and they highlight the advantages of population calcium imaging: excellent isolation of individual neurons and parallel sampling of local populations. Notably, cross-validation of our results is possible against data obtained with complementary methods. In particular, the receptive field sizes and scatter we observed are consistent with results from electrophysiological studies^{4,17–20}.

Shared and spanned subregions

The principal finding of our study is that RF subregions of layer 2/3 neurons in mouse visual cortex often precisely overlap with those of

other neurons distributed heterogeneously in the same local population. In addition, larger subregions seem to be composed of non-overlapping smaller subregions. These features limit the diversity of RFs and represent a fundamental principle of organization of the visual cortical microcircuit.

The subregions we observed are similar in size to the RFs of retinal ganglion cells²⁷. However, the subregions of cortical neurons are more elongated, with their aspect ratio ~ 1.6 on average, compared to ~ 1.2 for retinal ganglion cells²⁸. This likely reflects the integration of multiple afferents, possibly upstream of the layer 2/3 populations we have imaged. This integration may occur in the lateral geniculate nucleus, where neurons receive input from one to three retinal ganglion cells²⁹. Although RF properties in mouse visual cortex are quite similar to those observed in the lateral geniculate³⁰, we cannot exclude the possible contribution of further cortical integration upstream from the populations of neurons we imaged.

Neurons that shared subregions were not spatially clustered within visual cortex at the cellular scale, indicating that their dendritic fields did not necessarily need to overlap to produce identical subregions. This distributed processing of visual space stands in contrast to that in the retina, where individual mosaics of ganglion neurons are coordinated to sample visual space uniformly³¹. Similarly, neurons with subregions that spanned other subregions did not show a clear pattern of spatial organization within visual cortex. These spanning subregions may occur in neurons downstream in the circuit from those neurons expressing the subregions that they contain, suggesting the possibility of a processing hierarchy of RF subregions. Detailed circuit-tracing studies will be required to test this hypothesis.

Pairs of neurons that shared an RF subregion showed low correlations in activity, similar to their correlations with the rest of the population. This suggests that neurons sharing RF subregions remain independent, rather than simply serving as redundant elements in cortical circuitry. There are at least two possible schemes to consider. First, the neurons sharing an RF subregion may receive input that originates with the same small population of retinal ganglion cells. This naturally explains the high overlap but does not explain the low activity correlations. There are several potential sources of these low correlations. Low firing rates can limit the peak correlations (see **Supplementary Note 1**). RF subregions of the opposite (non-shared) sign are another mechanism that will limit correlations, by driving uncorrelated responses when the stimulus is not restricted to the shared RF subregion. Indeed, neurons that share a RF subregion of one sign have subregions of the opposite sign that overlap less (**Supplementary Fig. 8**). Similarly, neurons may be selective for different higher-order features of the visual stimulus that are not captured by the spike-triggered average^{32,33}. Finally, another potential source of the low correlations in this scenario is active decorrelation due to intracortical circuitry²¹.

Second, the two cells may receive input that originates from retinal ganglion cells (RGCs) that are of the same sign and the same region of visual space but of different types with different profiles of temporal response to the stimulus. The mouse retina is estimated to contain at least 20 different RGC types, each forming an independent mosaic^{34–36}. Furthermore, given the density of RGCs in the mouse retina and the surface area of visual cortex we imaged, we estimate that about 1,270 RGCs contributed to the region of visual space encoded in the portion of visual cortex we imaged (see **Supplementary Note 2**). Therefore, independent RGC input with the same sign encoding input for a very similar area of visual space is plausible and would naturally explain the low activity correlations. However, since each retinal mosaic is positioned independently, the

precise overlap seen in our data between different cell types would be expected to be uncommon. Hebbian mechanisms could alter receptive fields in the visual pathway and increase the functional overlap in primary visual cortex. It may be helpful to use population imaging to map RFs in developing visual cortex to determine whether the substantial overlap of subregions is a feature that is initially present or only emerges later owing to Hebbian mechanisms³⁷.

Spatially offset ensembles of ON and OFF RF subregions

We found the ON and OFF RF subregions of local populations of V1 neurons to be spatially offset in visual space. The origin of this phenomenon may lie in retinal ganglion cell mosaics (see **Supplementary Note 3**), but another source could also be responsible. In the optic tract of cats, retinal ganglion cell axons of the same type tend to travel together and form separate retinotopic maps that are slightly out of register with one another^{38,39}. A similar phenomenon may occur in mice. Although corrective mechanisms may exist to bring these maps back into register, such mechanisms will have to operate on pathways with very different firing patterns (ON and OFF cells for the same region of visual space). Thus, such mechanisms may fail to fully correct the registration error introduced by the fasciculation of axons of the same cell type.

Comparison with other species

Because many principles of visual cortex development and function are shared among mice, cats and primates^{40,41}, it will be interesting to determine whether the results presented here may also hold in these other species. In support of commonality, we found that clusters of ON and OFF subregions tended to be spatially segregated in visual space. This is reminiscent of the clustering of ON and OFF geniculate afferents in cat visual cortex⁴². The segregated processing of light and dark stimuli that we found at the local level may also have parallels in humans, given that human test subjects perform more poorly at detecting the interval between lines of opposite polarity than between lines of the same polarity, a phenomenon that has been interpreted as due to independent processing of light and dark stimuli^{43,44}. However, there are also aspects of the mouse visual system that differ from the visual systems of cats and primates (see **Supplementary Note 4**). Nevertheless, the principles we have uncovered are potentially applicable to intracortical connections with a low degree of divergence, which are found across a wide range of species.

Functional implications

Our results demonstrate that there is indeed robust functional micro-organization in mouse visual cortex that underlies the apparently weak local spatial organization of neurons with similar function previously described across rodent cortex⁴⁵. This result provides a useful framework with which to begin to dissect the cortical microcircuitry. Groups of neurons with shared or spanned RF subregions, despite precisely sharing a portion of their selectivity, act as independent, parallel units in the cortical circuitry.

This result demonstrates a mechanism for generating a high diversity of stimulus selectivities, despite low convergence: local scatter. The precisely shared RF subregions and spanned subregion sets indicate a low degree of convergence. If this low degree of convergence were combined with a low-scatter retinotopy, more neurons would have shared both ON and OFF subregions and local circuitry would have shown much less diversity. Using a high level of local RF scatter increases the diversity of stimulus selectivities. Thus, local RF scatter may be not a defect but rather a mechanism for generating diverse cortical computations.

Visual cortex circuitry can be considered in terms of discrete, self-contained circuits ('columns') or, alternatively, as a uniform network with smoothly varying stimulus selectivities. Our finding that the visual cortex contains groups of neurons that perform parallel processing of visual stimuli⁴⁶ of the same sign in the same area of visual space (at the expense of smooth local retinotopy) may lie somewhere between these two extremes. Groups of neurons can form discrete cell assemblies⁴⁷ by virtue of a shared subregion of the same sign and show a variety of stimulus selectivities due at least in part to their diversity of subregions of the opposite sign, thus forming a parallel processing unit for an area of visual space. However, these units are not mutually exclusive. Each neuron in a group may simultaneously participate in another group due to their subregions of the opposite sign, effectively participating in multiple cell assemblies⁴⁷. This clear evidence for an intermediate domain of discreteness raises many questions about models of cortical development and function. To address these questions, it will be important to determine the connectivity motifs of groups of neurons that share subregions and of neurons that show spanning subregions. This will ultimately require a combination of dense population imaging, as we have performed here, and new approaches for detailed cortical circuit analysis.

METHODS

Methods and any associated references are available in the online version of the paper at <http://www.nature.com/natureneuroscience/>.

Note: Supplementary information is available on the Nature Neuroscience website.

ACKNOWLEDGMENTS

We are grateful to B. Clark, M. Carandini, J. Cottam, M. Havenith, J. Jack, B. Judkewitz, P. Latham, M. London, A. Mathy, I. Smith and C. Wilms for discussions and for comments on the manuscript. This work was supported by a Human Frontier Science Program Long-Term Fellowship to S.L.S., and grants by the Wellcome Trust and the Gatsby Charitable Foundation to M.H.

AUTHOR CONTRIBUTIONS

S.L.S. and M.H. conceived the experiments. S.L.S. performed the experiments and analyzed the data. S.L.S. and M.H. interpreted the data and wrote the paper.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Published online at <http://www.nature.com/natureneuroscience/>.

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>.

- Hubel, D.H. & Wiesel, T.N. Receptive fields of single neurones in the cat's striate cortex. *J. Physiol. (Lond.)* **148**, 574–591 (1959).
- Smith, S.L. & Trachtenberg, J.T. Experience-dependent binocular competition in the visual cortex begins at eye opening. *Nat. Neurosci.* **10**, 370–375 (2007).
- Kalatsky, V.A. & Stryker, M.P. New paradigm for optical imaging: temporally encoded maps of intrinsic signal. *Neuron* **38**, 529–545 (2003).
- Wagor, E., Mangini, N.J. & Pearlman, A.L. Retinotopic organization of striate and extrastriate visual cortex in the mouse. *J. Comp. Neurol.* **193**, 187–202 (1980).
- Dräger, U.C. Receptive fields of single cells and topography in mouse visual cortex. *J. Comp. Neurol.* **160**, 269–290 (1975).
- Hubel, D.H. & Wiesel, T.N. Uniformity of monkey striate cortex: a parallel relationship between field size, scatter, and magnification factor. *J. Comp. Neurol.* **158**, 295–305 (1974).
- DeAngelis, G.C., Ghose, G.M., Ohzawa, I. & Freeman, R.D. Functional micro-organization of primary visual cortex: receptive field analysis of nearby neurons. *J. Neurosci.* **19**, 4046–4064 (1999).
- Reich, D.S., Mechler, F. & Victor, J.D. Independent and redundant information in nearby cortical neurons. *Science* **294**, 2566–2568 (2001).
- Yen, S.C., Baker, J. & Gray, C.M. Heterogeneity in the responses of adjacent neurons to natural stimuli in cat striate cortex. *J. Neurophysiol.* **97**, 1326–1341 (2007).
- Wässle, H., Boycott, B.B. & Illing, R.B. Morphology and mosaic of on- and off-beta cells in the cat retina and some functional considerations. *Proc. R. Soc. Lond. B Biol. Sci.* **212**, 177–195 (1981).
- Ringach, D.L. On the origin of the functional architecture of the cortex. *PLoS ONE* **2**, e251 (2007).
- Soodak, R.E. The retinal ganglion cell mosaic defines orientation columns in striate cortex. *Proc. Natl. Acad. Sci. USA* **84**, 3936–3940 (1987).
- Stosiek, C., Garaschuk, O., Holthoff, K. & Konnerth, A. In vivo two-photon calcium imaging of neuronal networks. *Proc. Natl. Acad. Sci. USA* **100**, 7319–7324 (2003).
- Ohki, K., Chung, S., Ch'ng, Y.H., Kara, P. & Reid, R.C. Functional imaging with cellular resolution reveals precise micro-architecture in visual cortex. *Nature* **433**, 597–603 (2005).
- Mata, M.L. & Ringach, D.L. Spatial overlap of ON and OFF subregions and its relation to response modulation ratio in macaque primary visual cortex. *J. Neurophysiol.* **93**, 919–928 (2005).
- Yeh, C.I., Xing, D. & Shapley, R.M. "Black" responses dominate macaque primary visual cortex V1. *J. Neurosci.* **29**, 11753–11760 (2009).
- Métin, C., Godement, P. & Imbert, M. The primary visual cortex in the mouse: receptive field properties and functional organization. *Exp. Brain Res.* **69**, 594–612 (1988).
- Mangini, N.J. & Pearlman, A.L. Laminar distribution of receptive field properties in the primary visual cortex of the mouse. *J. Comp. Neurol.* **193**, 203–222 (1980).
- Liu, B.H. *et al.* Visual receptive field structure of cortical inhibitory neurons revealed by two-photon imaging guided recording. *J. Neurosci.* **29**, 10520–10532 (2009).
- Niell, C.M. & Stryker, M.P. Highly selective receptive fields in mouse visual cortex. *J. Neurosci.* **28**, 7520–7536 (2008).
- Renart, A. *et al.* The asynchronous state in cortical circuits. *Science* **327**, 587–590 (2010).
- Ringach, D.L. Haphazard wiring of simple receptive fields and orientation columns in visual cortex. *J. Neurophysiol.* **92**, 468–476 (2004).
- Sohya, K., Kameyama, K., Yanagawa, Y., Obata, K. & Tsumoto, T. GABAergic neurons are less selective to stimulus orientation than excitatory neurons in layer II/III of visual cortex, as revealed by in vivo functional Ca2+ imaging in transgenic mice. *J. Neurosci.* **27**, 2145–2149 (2007).
- Mrsic-Flogel, T.D. *et al.* Homeostatic regulation of eye-specific responses in visual cortex during ocular dominance plasticity. *Neuron* **54**, 961–972 (2007).
- Gandhi, S.P., Yanagawa, Y. & Stryker, M.P. Delayed plasticity of inhibitory neurons in developing visual cortex. *Proc. Natl. Acad. Sci. USA* **105**, 16797–16802 (2008).
- Seung, H.S. Reading the book of memory: sparse sampling versus dense mapping of connectomes. *Neuron* **62**, 17–29 (2009).
- Balkema, G.W. Jr. & Pinto, L.H. Electrophysiology of retinal ganglion cells in the mouse: a study of a normally pigmented mouse and a congenic hypopigmentation mutant, pearl. *J. Neurophysiol.* **48**, 968–980 (1982).
- Hammond, P. Cat retinal ganglion cells: size and shape of receptive field centres. *J. Physiol. (Lond.)* **242**, 99–118 (1974).
- Chen, C. & Regehr, W.G. Developmental remodeling of the retinogeniculate synapse. *Neuron* **28**, 955–966 (2000).
- Grubb, M.S. & Thompson, I.D. Quantitative characterization of visual response properties in the mouse dorsal lateral geniculate nucleus. *J. Neurophysiol.* **90**, 3594–3607 (2003).
- Gauthier, J.L. *et al.* Receptive fields in primate retina are coordinated to sample visual space more uniformly. *PLoS Biol.* **7**, e1000063 (2009).
- Carandini, M. *et al.* Do we know what the early visual system does? *J. Neurosci.* **25**, 10577–10597 (2005).
- Olshausen, B.A. & Field, D.J. How close are we to understanding V1? *Neural Comput.* **17**, 1665–1699 (2005).
- Huberman, A.D. *et al.* Genetic identification of an On-Off direction-selective retinal ganglion cell subtype reveals a layer-specific subcortical map of posterior motion. *Neuron* **62**, 327–334 (2009).
- Huberman, A.D. *et al.* Architecture and activity-mediated refinement of axonal projections from a mosaic of genetically identified retinal ganglion cells. *Neuron* **59**, 425–438 (2008).
- Völgyi, B., Chheda, S. & Bloomfield, S.A. Tracer coupling patterns of the ganglion cell subtypes in the mouse retina. *J. Comp. Neurol.* **512**, 664–687 (2009).
- Song, S. & Abbott, L.F. Cortical development and remapping through spike timing-dependent plasticity. *Neuron* **32**, 339–350 (2001).
- Torrealba, F., Guillery, R.W., Eysel, U., Polley, E.H. & Mason, C.A. Studies of retinal representations within the cat's optic tract. *J. Comp. Neurol.* **211**, 377–396 (1982).
- Torrealba, F., Guillery, R.W., Polley, E.H. & Mason, C.A. A demonstration of several independent, partially overlapping, retinotopic maps in the optic tract of the cat. *Brain Res.* **219**, 428–432 (1981).
- Nauhaus, I., Benucci, A., Carandini, M. & Ringach, D.L. Neuronal selectivity and local map structure in visual cortex. *Neuron* **57**, 673–679 (2008).
- Gordon, J.A. Cellular mechanisms of visual cortical plasticity: a game of cat and mouse. *Learn. Mem.* **4**, 245–261 (1997).
- Jin, J.Z. *et al.* On and off domains of geniculate afferents in cat primary visual cortex. *Nat. Neurosci.* **11**, 88–94 (2008).
- Levi, D.M., Jiang, B.C. & Klein, S.A. Spatial interval discrimination with blurred lines: black and white are separate but not equal at multiple spatial scales. *Vision Res.* **30**, 1735–1750 (1990).
- Levi, D.M. & Westheimer, G. Spatial-interval discrimination in the human fovea: what delimits the interval? *J. Opt. Soc. Am.* **4**, 1304–1313 (1987).
- Ohki, K. & Reid, R.C. Specificity and randomness in the visual cortex. *Curr. Opin. Neurobiol.* **17**, 401–407 (2007).
- Gao, E., DeAngelis, G.C. & Burkhalter, A. Parallel input channels to mouse primary visual cortex. *J. Neurosci.* **30**, 5912–5926 (2010).
- Hebb, D.O. *The Organization of Behavior: A Neuropsychological Theory* (Wiley, 1949).

ONLINE METHODS

Preparation. All experiments were carried out in accordance with UK Home Office regulations. Adult C57BL/6 mice (1.5–3 months of age) were anesthetized with isoflurane (5% for induction, 1–3% for surgery) and chlorprothixene (0.5–2 mg per kilogram body weight). During imaging, the isoflurane was decreased to 0–0.5%. A 2–3 mm diameter craniotomy was opened over monocular visual cortex. The dura mater was left intact. Bolus loading of neurons with calcium dye was performed using a standard protocol^{13,14,48}. A 50- μ g aliquot of Oregon Green-BAPTA (1,2-bis-(*o*-aminophenoxy)ethane-*N,N,N',N'*-tetra-acetic acid)-1 or Fluo-4 (Invitrogen) was dissolved in a 20% Pluronic-127 (Invitrogen) solution in DMSO. After vortexing, this was diluted tenfold with a pH 7.4 solution of 150 mM NaCl, 2.5 mM KCl, 10 mM HEPES and 50 μ M sulforhodamine 101 to counterstain astrocytes⁴⁹. After filtering the solution, we guided a patch pipette (3–7 M Ω) under two-photon microscopy to a depth of 150–250 μ m. Pressure of 300–600 mbar was applied for 1–3 min to eject dye solution from the pipette. This stained a spherical volume of 300–400 μ m in diameter. We imaged 434 neurons in six mice and obtained RFs from 189 of these cells (44%).

Imaging. A custom two-photon microscope with a $\times 16$, 0.8 numerical aperture (Nikon) water immersion objective and a large aperture collection pathway with low-noise photomultiplier tubes (hand-selected 3896 and 7422-40P photomultiplier tubes, Hamamatsu) was used for imaging. Images were acquired using ScanImage in conjunction with MATLAB at 15.6 frames per second with 256×128 pixels resolution, covering an area of 230×230 μ m.

Electrophysiology. To calibrate the deconvolution routine, on-cell recordings were performed during imaging and stimulus presentation. Standard techniques⁵⁰ were used to visually guide the pipette (3–5 M Ω), which contained HEPES-buffered artificial cerebrospinal fluid and 25–50 μ M Alexa 594 (Invitrogen), to the target neuron. Recordings were obtained in current-clamp mode with no current injection. Signals were filtered at 3–5 kHz and acquired at 10–50 kHz.

Visual stimulation. Routines written in MATLAB using the Psychophysics Toolbox (Brainard and Pelli) extensions controlled the visual stimulus presentation. Sparse-noise visual stimuli consisted of black (2 cd m⁻²) and white (86 cd m⁻²) dots on a gray (40 cd m⁻²) background¹⁵. Dots ranged in size from 1.3–8.0° in diameter. Adjacent frames had no overlap in dot locations so that all pixel transitions were to or from gray, never from black to white or vice versa. The visual stimulus LCD screen (ESAW 7 inch VGA TFT, set at 1024 $\times$ 768 resolution and 60 Hz refresh rate) was shrouded with a cone up to the eye of the mouse to prevent contamination of the imaging pathway with light from the visual stimulus. The visual stimulus extended from +20°–+124° in azimuth (in one experiment, the span was +20°–+73°) and from –10°–+42° in elevation. Stimulus frames (128 $\times$ 128 pixels) were smoothly interpolated so that 1 stimulus frame pixel was equivalent to 0.72°² in visual space. Between 4,000 and 6,000 stimulus frames were used in each experiment. Total mapping time was 30–45 min.

Analysis and statistics. In the calcium imaging movies of visual cortex during stimulus presentation, regions of interest were detected using a local pixel-wise neighborhood cross-correlation (Supplementary Fig. 7). For a given movie, the time course for each pixel was compared to those of its eight neighbors. The average peak cross-correlation value was then used to form a local cross-correlation peak image. This image was thresholded using a locally adaptive filter to obtain regions of interest. Subsequently, morphological filters were used to identify somata.

After filtering of the fluorescence time courses with a locally adaptive Wiener filter, a deconvolution-based algorithm generated estimated spike rates. The deconvolution kernel was an instantaneous rise followed by an exponential decay. The time course of the exponential decay was selected on the basis of the electrophysiology data to maximize the match between the real and estimated spiking. Once optimized, the same kernel ($\tau_{\text{decay}} = 0.64$ s) was used for all time courses. The resulting trace was then thresholded to remove noise; this threshold was adaptive in that it was adjusted on the basis of the s.d. and skewness of the deconvolved traces. When the threshold was adjusted manually for each cell individually, the resulting cross-correlations between recorded spike trains and estimated spike trains did not change noticeably (automatic noise threshold cross-correlation, 0.81 ± 0.02 ; manual noise threshold cross-correlation, 0.84 ± 0.02).

When the noise threshold was selected on the basis of all but one data set, and then tested on the excluded data set, the estimated spike trains again did not change noticeably (cross-correlation, 0.81 ± 0.02). Detection rates (Fig. 1d) were calculated as the likelihood that any given, single imaging frame event will be detected. This detection rate does not factor in the accuracy of event amplitude estimation (number of spikes). The cross-correlation measure does depend on the accuracy of the event amplitude estimation but is insensitive to changes in scale. We restricted our analysis to measures that were also insensitive to scale: namely, spike-triggered averages and cross-correlations.

These estimated spike-rate time courses were used for reverse correlation with the visual stimulus and for determining cross-correlations between neurons. Raw fluorescence traces were not used for computing cross-correlations because of concerns that they can yield artificially high correlations (due to a variety of causes, including slow decay time courses after events and small fluctuations in laser power). Indeed, when cross-correlations were computed on raw fluorescence traces, the resulting cross-correlation values were higher than that obtained from deconvolved traces (Supplementary Fig. 8).

We did not explicitly differentiate between excitatory and inhibitory neurons. However, our deconvolution scheme relies on the well-characterized calcium signal in response to action potentials, and the action potential-associated calcium signals in interneurons are typically markedly smaller than those of principal neurons, owing to the robust calcium buffering mechanisms specific to interneurons. Together with the higher prevalence of pyramidal neurons compared to interneurons in mammalian cortex, it is therefore likely that the population of neurons reported here is dominated by pyramidal neurons.

Reverse correlation was computed using the delay that yielded the highest signal-to-noise ratio in the RF maps across all neurons (mean, 255 ms; range, 200–300 ms). The same delay parameter was used for all neurons within a population. Estimated spike-triggered averages were thresholded at ± 4 z scores. To further decrease the noise in the RF maps, a morphological filter excluded small features ($\leq 2.6^\circ$ in radius). Negative z -score areas (where visual stimuli suppressed spiking) were observed but were rare and typically weak. For simplicity, we focused solely on the positive z -score areas for this analysis. For clarity, interpolated RF contours are shown in some figures. All analysis was performed on uninterpolated (pixel-wise) binary RFs.

The overlap index metric was calculated as the proportion of the smaller of two subregions that overlapped the larger subregion. Shared subregions were defined as having $\geq 90\%$ overlap, centers $\leq 1.5^\circ$ from each other and $\leq 25\%$ difference in area. Spanned subregion sets were defined as an ensemble of two or more subregions with $\leq 25\%$ overlap and another subregion that overlapped the ensemble $\geq 75\%$. To measure the incidence of spanned sets, we calculated the number of unique span sets in the population, normalized by the number of subregions; a span set was defined as a set of non-overlapping subregions and a corresponding spanning subregion. The main results of this study were not acutely sensitive to these definitions. We also compared an alternative measure of subregion overlap to the overlap metric described above. This alternative was a subregion correlation (SC) metric that was calculated using the formula

$$SC = 1 - \frac{\sum_{\text{all pixels}} |srA - srB|}{\sum_{\text{all pixels}} srA + \sum_{\text{all pixels}} srB}$$

where srA and srB are binary images of two subregions with backgrounds of zeros. For this metric, 0 indicates completely non-overlapping subregions, increasing values represent greater amounts of subregion overlap and a value of 1 indicates that the subregions are identical. The results obtained using the overlap metric with the thresholds described above and those obtained using the subregion correlation metric with a threshold of 0.85 were indistinguishable. Thus, the results presented in this study do not depend on a specific quantification of subregion overlap.

We also computed image correlations using another metric. The two images were each converted into one-dimensional series, similar to two time series, of pixels, A and B . The correlation R was computed as

$$R = \frac{E[(A - \mu_A)(B - \mu_B)]}{\sigma_A \sigma_B}$$

where μ_X and σ_X are the mean and s.d. of series X , respectively, and E is the expected value operator. When this metric was used to compare z -scored maps, the maps were first masked by a two-dimensional Gaussian mask with an s.d. equal to the longest radius of the RF subregion. This minimally affected the z -scored maps near the RF subregion and flattened out the background regions to values near 0. This approach was taken to ensure that the correlation coefficients calculated were dominated by correlations in or near the RF subregion and not by the background noise.

Span sets were found using a tree search algorithm. Neurons with similar, but not necessarily shared, subregions could increase the number of span sets. For example, take the case in which neurons A, B and C have similar subregions; likewise for neurons D, E, and F; and neurons G and H have similar subregions, which span the previous ones. Then the spanning subregions sets would be ADG, BDG, CDG, AEG, ... CFH: a total of 18 unique sets for 8 neurons. Note that, in practice, not all shared subregions will degenerately contribute to the same spanning subregion count; some shared subregions will vary just enough to exclude them and vice versa for subregions in the spanning subregion count being excluded from the shared subregions. Because shared RF subregions can contribute to high-spanning set counts and thus confound a statistical test of the spanning RF subregions, we relaxed the thresholds for the spanning RF subregion test to the point where the statistical significance for shared RF subregions breaks down. Thus, any remaining difference between observed and repositioned subregions was due to the feature of spanning RF subregions. The potential for high counts in this analysis was mitigated by using the non-parametric, paired, two-sided Wilcoxon signed-rank test for determining statistical significance.

To test whether the observed ensembles of subregions were significantly different from other random spatial configurations, we compared the ensembles to randomly repositioned versions. To construct these, observed subregions were randomly repositioned in visual space. The repositioning did not change subregion shapes or orientation. The repositioning preserved the scatter and spatial spread (**Supplementary Fig. 9**) of subregion ensembles in visual space, as observed in the original data. To obtain average values for the randomly repositioned versions of ensembles, the repositioning was repeated 1,000 times.

The clustering of ON and OFF subregions was quantified by subtracting the sum of the OFF subregions from the sum of ON subregions. In this computation, highly overlapping ON and OFF subregions will result in an image that is on average zero, and any clustering will result in positive and negative peaks. The sum of absolute values of all pixels was then used as a measure of clustering. To test significance, the subregions were randomly reassigned as ON or OFF and the clustering was recalculated. After 1,000 iterations, the mean of the clustering values for the reassigned data was used to normalize the observed clustering measures.

Unless otherwise noted, all measurements are expressed as mean $\pm$ s.e.m.

48. Garaschuk, O., Milos, R.I. & Konnerth, A. Targeted bulk-loading of fluorescent indicators for two-photon brain imaging *in vivo*. *Nat. Protoc.* **1**, 380–386 (2006).
49. Nimmerjahn, A., Kirchhoff, F., Kerr, J.N. & Helmchen, F. Sulforhodamine 101 as a specific marker of astroglia in the neocortex *in vivo*. *Nat. Methods* **1**, 31–37 (2004).
50. Kitamura, K., Judkewitz, B., Kano, M., Denk, W. & Häusser, M. Targeted patch-clamp recordings and single-cell electroporation of unlabeled neurons *in vivo*. *Nat. Methods* **5**, 61–67 (2008).